



Receipt number	662-06-E-4231
Study number	94231

FINAL REPORT

A 96-hour Acute Toxicity Study of 13F-OLE with Medaka

September 19, 2007

Kurume Laboratory
Chemicals Evaluation and Research Institute, Japan

STATEMENT

Kurume Laboratory
Chemicals Evaluation and
Research Institute, Japan

Sponsor DAIKIN INDUSTRIES, LTD.

Title A 96-hour Acute Toxicity Study of 13F-OLE with Medaka

Study number 94231

I, the undersigned, hereby declare that this report provides a correct English translation of the Final Report (Study No. 94231, issued on September 19, 2007)

Date *October 27, 2009*

Study Director *Takeshi Matsuura*
Takeshi Matsuura

CONTENTS

	Page
Title	1
Sponsor.....	1
Test facility.....	1
Objective	1
Test method.....	1
Applied GLP	1
Dates.....	2
Storage of test item, raw data, etc.	2
Personnel	3
Approval of final report	3
SUMMARY.....	4
1. Test item	6
2. Test sample.....	7
3. Test materials and methods	8
4. Results and discussion	12
5. Factors that affected reliability of test results	14

Tables

Table 1	Cumulative mortality
Table 2	Observed abnormal response
Table 3-1	Dissolved oxygen concentration of test solutions
Table 3-2	pH of test solutions
Table 3-3	Temperature of test solutions
Table 4	LC ₅₀ to Medaka
Appendix 1	Chemical characteristics of dilution water
Appendix 2	Analytical method and measured concentration of test item
Appendix 3	Calibration curve and chromatogram
Appendix 4	Solubility in dilution water
Additional data	Results of preliminary studies

Title	A 96-hour Acute Toxicity Study of 13F-OLE with Medaka
Sponsor	DAIKIN INDUSTRIES, LTD. 1-1, Nishi-Hitotsuya, Settsu, Osaka 566-8585, Japan
Test facility	Kurume Laboratory Chemicals Evaluation and Research Institute, Japan 3-2-7 Miyanojin, Kurume-shi, Fukuoka 839-0801, Japan
Objective	The purpose of this study is to determine the acute toxicity of 13F-OLE on fish.
Test method	This study was performed according to the following test methods and guidance document. <ol style="list-style-type: none">(1) Fish, Acute Toxicity Test stipulated in the "Testing Methods for New Chemical Substances" (November 21, 2003; No. 1121002, Pharmaceutical and Food Safety Bureau, Ministry of Health, Labour and Welfare; November 13, 2003, No. 2, Manufacturing Industries Bureau, Ministry of Economy, Trade and Industry; No. 031121002, Environmental Policy Bureau, Ministry of the Environment)(2) OECD Guidelines for Testing of Chemicals, Section 2 : Effects on Biotic Systems, 203 Fish, Acute Toxicity Test (Guideline 203, 1992)(3) OECD Guidance Document No.23 "Guidance Document on Aquatic Toxicity Testing of Difficult Substance and Mixtures" (September 2000)
Applied GLP	This study was conducted in compliance with the following GLP principles: <ol style="list-style-type: none">(1) "Standard Concerning Testing Facility Relating to New Chemical Substances" (November 21, 2003; No. 1121003, Pharmaceutical and Food Safety Bureau, Ministry of Health, Labour and Welfare; November 17, 2003, No. 3, Manufacturing Industries Bureau, Ministry of Economy, Trade and Industry; No. 031121004, Environmental Policy Bureau, Ministry of the Environment)(2) "OECD Principles of Good Laboratory Practice (November 26, 1997, ENV/MC/CHEM(98)17)"

Dates

Study initiation date	August 15, 2007
Experimental starting date	August 20, 2007
Solubility study starting date	August 20, 2007
Bioassay starting date	August 20, 2007
Experimental completion date	August 24, 2007
Solubility study completion date	August 22, 2007
Bioassay completion date	August 24, 2007
Study completion date	September 19, 2007

Storage of test item, raw data, etc.

(1) Test item

The test sample^{*1} will be sealed in a storage vessel and stored in archives in this laboratory for ten years after the receipt of notice specified under Clause 1 or Clause 2 in Article 4, Clause 2 or Clause 3 or Clause 8 in Article 4-2, and Clause 2 in Article 5-4 or Clause 2 in Article 24 or Clause 2 in Article 25-3 of "Law Concerning Examination and Regulation of Manufacture, etc. of Chemical Substances". If it is not stable for the storage period, it is stored as long while it is kept stable. Treatment of the sample after the storage period will be discussed with sponsor.

*1 It will be stored as the common sample for storage of these studies (Study Nos. 94229, 94230 and 94231).

(2) Raw data and materials

Raw data, the study plan, documents concerning the study presented by the sponsor, the final report and necessary materials will be stored in archives in this laboratory for ten years after the receipt of the notice specified under Clause 1 or Clause 2 in Article 4, Clause 2 or Clause 3 or Clause 8 in Article 4-2, and Clause 2 in Article 5-4 or Clause 2 in Article 24 or Clause 2 in Article 25-3 of "Law Concerning Examination and Regulation of Manufacture, etc. of Chemical Substances". Treatment of raw data and materials, etc. after the storage period will be discussed with sponsor.

Personnel

Study Director : Takeshi Matsuura
Section 4 (Eco-toxicity test area)

Study personal

Biology : Takeshi Matsuura, Miyuki Narasaki,
Jun Sugimura, Makiko Anai,
Mika Ono, Mariko Tanaka,
Yumika Hiroo

Analytical chemistry : Junji Azuma, Tomoyo Kagamiyama,
Hiroko Sugimoto, Hitomi Inoue,
Hitomi Fukui, Hiromi Andoh

Approval of final report

Study Director Date September 19, 2007

Signature Signed in original
Takeshi Matsuura

SUMMARY

Title

A 96-hour Acute Toxicity Study of 13F-OLE with Medaka

Test conditions

(1) Test item	13F-OLE
(2) Test organism	Medaka (<i>Oryzias latipes</i>)
(3) Exposure duration	96 hours
(4) Test concentration	Middle layer of suspended solution (nominal concentration: 100 mg/L), and control
(5) Replicate	Two replicates/test level
(6) Number of organism	Ten fish / test level (five fish / test vessel)
(7) Dilution water	Dechlorinated tap water
(8) Type of test	Semi-static regime (renewal at every 24 hours) with closed system
(9) Preparation of test solution	The test sample and dilution water were mixed to produce 100 mg/L (nominal concentration), and they were stirred under closed system for approximately 48 hours. After settlement for approximately 1 hour, test solution was prepared by taking out from the middle layer.
(10) Volume of test solution	Approximately 6 L / test level (Approximately 3 L / test vessel)
(11) Temperature of test solutions	24±1°C
(12) Irradiation condition	Artificial light of white fluorescent lamp, 16-hour light / 8-hour dark
(13) Feeding	No feeding
(14) Aeration	No aeration
(15) Analysis of concentration of test item in test solution	GC-MS analysis (at the start of the exposure, before and after the renewal, and the end of the exposure)

Results

(1) Solubility of test item in dilution water (24±1°C)	0.101 mg/L
(2) Concentration of test item in test solution (Percentage of concentration at preparation)	
At the start of the exposure and after the renewal	0.0805 to 0.329 mg/L
Before the renewal and at the end of the exposure	0.0666 to 0.168 mg/L (41.2 to 95.4%)
(3) 96-hour LC ₅₀ (Median Lethal Concentration)	> 0.117 mg/L

[The values of (3) is based on geometric mean of the measured concentrations.]

Conclusion

This study was conducted as a limit test at the concentration around the solubility of the test item in dilution water to confirm the effect on the test organisms. It was concluded that the test item has no acute toxicity to the test organisms at the concentration around the solubility, since the measured concentrations of the test solutions were around the solubility in dilution water at the preparation and no effect on the test organisms was observed under the test condition.

1. Test item


In this report, 13F-OLE has the following name etc.

1.1 Chemical name*²

3,3,4,4,5,5,6,6,7,7,8,8,8-tridecafluoro-octa-1-ene

1.2 Chemical structure etc.*²

Structural formula



Molecular formula $\text{C}_8\text{H}_3\text{F}_{13}$

Molecular weight 346.09

CAS Number 25291-17-2

*² Information supplied by the sponsor

2. Test sample

2.1 Supplier and lot number^{*2}

Supplier	DAIKIN INDUSTRIES, LTD.
Lot number	061122HM

2.2 Purity^{*2}

Test item	99.8%
Impurity	Unknown constituent component 0.2%

2.3 Confirmation of test item

It was confirmed that infrared (IR) spectrum of the test item provided by the sponsor coincided with IR spectrum analyzed in this laboratory.

2.4 Physicochemical properties^{*2}

Appearance at normal temperature	Colorless and clear liquid	
Boiling point	106°C (760 mmHg)	
Density	1.560 g/cm ³ (20°C)	
Solubility	Water	Insoluble
	Dimethylsulfoxide	Insoluble
	Acetone	Soluble (fully miscible)

^{*2} Information supplied by the sponsor

2.5 Storage condition and confirmation of stability at storage condition

Storage condition	Dark storage place at room temperature
Confirmation of stability	The stability of the test item during the test period was confirmed by no alteration in the IR spectra of the test item before the experimental start and after the experimental completion.

3. Test materials and methods

3.1 Test organism

(1) Species

Medaka (*Oryzias latipes*)

(2) Reason for selection of species

This species is recommended in the test guidelines.

(3) Size

Total length 2.3±1.2 cm

Size of test organism was applied the regulated size set to test method (1).

(4) Supplier

Kurume Laboratory, Chemicals Evaluation and Research Institute, Japan
3-2-7 Miyanojin, Kurume-shi, Fukuoka 839-0801, Japan

(5) Acclimation

Medaka hatched out on March 27, 2007 were acclimated for 27 days by flow-through condition under the same water quality (dechlorinated tap water), temperature (24±1°C), light and dark period (16-hour light / 8-hour dark) as test condition (on July 24 to August 20, 2007). The mortality was 0% during the 7 days before the start of the exposure. The test organism at the start of the exposure was 4-month-old fish. The test organisms were not treated with a medicament for external disinfection. The test organisms were fed the feed mixture for carp (2C), and not fed for 24 hours before the start of the exposure. Dissolved oxygen concentration in breeding water during acclimation was kept not less than 80% of air saturation value. A 96-hour acute toxicity test of CuSO₄ 5H₂O (Reagent chemical, Wako Pure Chemical Industries, Ltd.) to confirm reproducibility of the test system was carried out on May 21 to May 25, 2007 and the 96-hour LC₅₀ was 0.445 mg/L. This value was within the stipulated range (mean ± 2S.D. : 0.124 to 0.978 mg/L) [mean ± S.D. : 0.551 ± 0.214 mg/L (n=38)] to background data in this laboratory. All of the values shown above for the reference substance were converted into CuSO₄ value.

(6) Allocation to the test groups

Medaka were allocated at random to each test group.

3.2 Dilution water

Dechlorinated tap water, aerated sufficiently and controlled temperature, was used. Some chemical characteristics of the dilution water measured regularly are listed in Appendix 1.

3.3 Test apparatus and equipment

(1) Test apparatus

Test vessel : 3 L Glass tank (diameter: 16 cm, depth: 17 cm)

The test vessels were covered with a glass lid in order to prevent dust, and volatilization of the test solution and closed without headspace.

(2) Test equipment

Water bath : Plastic tank

Warming / cooling unit (Type HCA 250, Sato craft)

3.4 Test conditions

(1) Conditions of exposure

(a) Type of test

The test organisms were exposed to the test solution containing the test item.

The test solutions were renewed at every 24 hours, as closed semi-static regime.

(b) Exposure duration

96 hours

(c) Test concentration

Based on the results of the preliminary studies, it was expected that the test solution at around the solubility in dilution water would have no effect on the test organisms. Therefore, the definitive study was conducted as the limit test with suspended solution which was prepared by taking out from the middle layer of approximately 48-hour mixed solution (nominal concentration: 100 mg/L). The results of the preliminary studies are shown in Additional data.

(d) Control

The dilution water without the test item, which was treated in the same stirring manner as the test solution, was used as the control.

(e) Replicates

Two replicates / test level

- (f) Number of organism
Ten fish / test level (five fish / test vessel)
 - (g) Volume of test solution
Approximately 6 L / test level (approximately 3 L / test vessel)
- (2) Conditions of test environment
- (a) Water temperature
24±1°C
 - (b) Dissolved oxygen concentration
The test was performed in the condition where dissolved oxygen concentration was at least 60% or more of the saturated concentration at the water temperature. Aeration was not used during the exposure.
 - (c) pH
The test was performed without adjusting pH.
 - (d) Irradiation condition
Artificial light of white fluorescent lamp, 16-hour light/8-hour dark
 - (e) Feeding
Test organisms were not fed during the exposure.

3.5 Preparation of test solution

No correction with purity was done for the preparation of the test concentration. The test sample was employed in terms of volume using the density [1.560 g/cm³ (20°C)] for the preparation of test solution.

After the test sample was added into the dilution water filled in Erlenmeyer flask with micro volumeter (Eppendorf Co., Ltd) to produce 100 mg/L as nominal concentration, the flask was immediately sealed with a plug not to produce head space. The solution was gently stirred by magnetic stirrer for approximately 48 hour. After cease of stirring, the solution was settled for approximately 1 hour under 24±1°C and then test solution was prepared by taking out from the middle layer of the settled solution. The prepared test solution was immediately divided into each test vessel and covered with glass lid not to produce head space.

3.6 Observation and measurements

(1) Observation of test organisms

Mortality and visible abnormality were observed at 3, 24, 48, 72 and 96 hours after the start of the exposure. A fish was considered as dead if the observable motion (motion of mouth and opercula etc.) were not observed and touching of the caudal peduncle with glass rod produced no reaction.

(2) Total length and body weight of test organism

The test organisms in the control group were used for measuring total length and body weight after the end of the exposure.

(3) Appearance of test solution

Appearance of the test solutions were observed at the start of the exposure and before the renewal (after 24 hours).

(4) Condition of test solutions

Dissolved oxygen concentration, pH and temperature of the test solutions were measured at the start of the exposure, before and after the renewal and the end of the exposure. At the start of exposure and after the renewal (at the preparation), another solution sampled from the container for preparation was used for the measurement. Before the renewal and the end of exposure, the measurement was carried out for one test vessel in each level. The dissolved oxygen concentration measurements were carried out with an oxygen meter (YSI Incorporated., YSI Model 58). The pH measurements were carried out with a portable pH meter (DKK-TOA, Model HM-21P). The temperature measurements were carried out with a calibrated red alcohol thermometer of glass stick type.

(5) Concentration of test item in test solution

The concentration of the test item in the test solutions was measured at the start of the exposure, before and after the renewal and the end of the exposure. At the start of the exposure and after the renewal (at the preparation) another solution sampled from the container for preparation was used for analysis. Before the renewal and the end of the exposure, the test solution for analysis was taken out with equal volume from the middle layer of the test solution in test vessels in each test level, and mixed. The concentration of the test item was analyzed by gas chromatography - mass spectrometry (GC-MS). Analytical method and measured concentration of test item are shown in Appendix 2, and analytical calibration curve and chromatograms are shown in Appendix 3.

(6) Solubility of test item in dilution water

Since the solubility of the test item was expected less than 100 mg/L, it was measured concurrently with the definitive study. The detail and the result of the measurement of the solubility are shown in Appendix 4.

3.7 Calculating method of LC₅₀^{*3}

The LC₅₀ values were estimated as “> the test concentration” since no less than 50% of mortality was observed in the exposure level.

The results of the study were estimated based on a geometric mean of the measured concentrations as the test concentration.

*3 LC₅₀ (Median Lethal Concentration): The test item concentration at which 50% of the test organisms causes mortality during the exposure.

3.8 Validity of the test

- (1) The mortality in the control should not exceed 10%.
- (2) Dissolved oxygen concentration must be at least 60% of the air saturation value at the water temperature in the test during exposure.

3.9 Treatment of numerical values

Values were rounded off in accordance with JIS Z 8401 rule B, 1999.
(JIS; Japanese Industrial Standards)

4. Results and disssion

4.1 Mortality

The mortality during the exposure was not observed in the exposure level. Cumulative mortality of each exposure period was shown in Table 1. The mortality in the control was not observed, which meets the criterion for the validity of the test (i.e. not more than 10%).

4.2 Observed abnormal response

No abnormal responses were observed in the control.

The results of observation were based on the comparison with the control organisms. No abnormal responses were observed in the exposure level during exposure. The abnormal responses observed during the exposure are shown in Table 2.

4.3 Size of test organism [Mean ± Standard deviation (n=10)]

Total length	2.4±0.11 cm
Body weight	0.11±0.013 g

4.4 Observation and measurement of test solution

(1) Appearance of test solution

The test solutions were clear and colorless at the start of the exposure and before the renewal.

(2) Condition of test solutions

The measured values of dissolved oxygen concentration, pH and temperature during the exposure ranged from 6.8 to 8.4 mg/L, 7.5 to 7.8 and 24.0 to 24.7°C, respectively. Conditions of the test solutions are shown in Tables 3-1, 3-2 and 3-3. The measured values of dissolved oxygen concentration met the criterion for the study validity (at least 60% or more of saturate concentration*⁴ at the water temperature).

*4 Saturated dissolved oxygen concentration (23 to 25°C): 8.39 to 8.11 mg/L
(JIS K 0102, 1998)

(3) Concentration of test item in test solution

The measured concentrations of the test item in the test solution were 0.0805 to 0.329 mg/L at the start of the exposure and after the renewal. Before the renewal and at the end of the exposure, they were 0.0666 to 0.168 mg/L which were 41.2 to 95.4% of the concentration at the preparation. The results of the measured concentrations of the test item are shown in Appendix 2.

4.5 LC₅₀

Both of the 48 and 96-hour LC₅₀s of the test item to Medaka were >0.117 mg/L.
The LC₅₀s at each time are shown in Table 4.

4.6 Discussion

This study was conducted as a limit test in order to confirm the effect of the test item on the test organisms at the concentration around the solubility of the test item in dilution water. As a result, the measured concentration of the test solution at the preparation was almost the same concentration as the solubility in the dilution water. Although, it decreased at 24 hours after the preparation. It is thought that the definitive study was appropriate for the test at the concentration around the solubility since the test was performed using semi-static replacement regime (renewal at 24 hours) to maintain the test concentration. No adverse effect was found under the condition in the definitive study, therefore, it was concluded that the test item had no adverse acute effect on the test organisms at around the solubility in dilution water

5. Factors that affected reliability of test results

There were no factors which might have affected the reliability of the test.

Table 1 Cumulative mortality

Measured concentration *5 (mg/L)	Cumulative mortality (%)				
	3 hours	24 hours	48 hours	72 hours	96 hours
Control	0	0	0	0	0
0.117	0	0	0	0	0

*5 Geometric mean of the measured concentrations

(The followings are expressed as measured concentration)

Table 2 Observed abnormal response

Measured concentration (mg/L)	Result of observation (Left column: Number of affected fish/Total survival number, Right column: Symptom detail)									
	3 hours		24 hours		48 hours		72 hours		96 hours	
	Control	0/10	N	0/10	N	0/10	N	0/10	N	0/10
0.117	0/10	N	0/10	N	0/10	N	0/10	N	0/10	N

N : Normal (No abnormal response)

Table 3-1 Dissolved oxygen concentration of test solutions

Measured concentration (mg/L)	0 hour	24 hours		48 hours		72 hours		96 hours
	At the start	Before the renewal	After the renewal	Before the renewal	After the renewal	Before the renewal	After the renewal	At the end
Control	8.2	7.8	8.4	8.0	8.2	7.1	8.1	7.0
0.117	8.0	7.4	8.1	7.9	8.2	6.9	8.0	6.8

Unit : mg/L

Table 3-2 pH of test solutions

Measured concentration (mg/L)	0 hour	24 hours		48 hours		72 hours		96 hours
	At the start	Before the renewal	After the renewal	Before the renewal	After the renewal	Before the renewal	After the renewal	At the end
Control	7.8	7.5	7.7	7.5	7.8	7.6	7.7	7.5
0.117	7.8	7.5	7.8	7.5	7.8	7.6	7.8	7.5

Table 3-3 Temperature of test solutions

Measured concentration (mg/L)	0 hour	24 hours		48 hours		72 hours		96 hours
	At the start	Before the renewal	After the renewal	Before the renewal	After the renewal	Before the renewal	After the renewal	At the end
Control	24.2	24.0	24.1	24.1	24.3	24.0	24.6	24.1
0.117	24.7	24.0	24.1	24.0	24.0	24.1	24.5	24.0

Unit: °C

Table 4 LC₅₀ to Medaka

Exposure duration	LC ₅₀ (mg/L)	95% confidence interval (mg/L) (Slope of the dose-response curve)	Statistical procedure used for determination of LC ₅₀
24-hour	>0.117	- (-)	-
48-hour	>0.117	- (-)	-
72-hour	>0.117	- (-)	-
96-hour	>0.117	- (-)	-

- : Not obtained

Appendix 1

Chemical characteristics of dilution water

Chemical characteristics of dilution water (Sampling on July 2, 2007)

Parameter	Unit	Results	Lower limit of determination
Total hardness (as CaCO ₃)	mg/L	37.0	0.1
Suspended solid	mg/L	< 1	1
pH	—	7.7 (24°C)	—
Total organic carbon	mg/L	< 0.1	0.1
Chemical oxygen demand	mg/L	< 0.5	0.5
Residual chlorine	mg/L	< 0.02	0.02
Ammonium ion	mg/L	0.02	0.01
Total cyan	mg/L	< 0.01	0.01
Alkalinity	mg/L	29	1
Electric conductivity	mS/m	15.5	—
Organic phosphorous	mg/L	< 0.1	0.1
Alkylmercury	mg/L	< 0.0005	0.0005
Mercury	mg/L	< 0.0005	0.0005
Cadmium	mg/L	< 0.001	0.001
Chromium (VI)	mg/L	< 0.02	0.02
Lead	mg/L	< 0.005	0.005
Arsenic	mg/L	< 0.001	0.001
Boron	mg/L	0.04	0.02
Fluorine	mg/L	0.1	0.1
Iron	mg/L	< 0.01	0.01
Copper	mg/L	< 0.005	0.005
Cobalt	mg/L	< 0.001	0.001
Manganese	mg/L	< 0.01	0.01
Zinc	mg/L	< 0.005	0.005
Aluminum	mg/L	0.051	0.001
Nickel	mg/L	< 0.001	0.001
Silver	mg/L	< 0.0001	0.0001
Sulfate ion	mg/L	12.9	0.1
Chloride ion	mg/L	15	1
Sodium	mg/L	13.1	0.01
Potassium	mg/L	3.6	0.01
Calcium	mg/L	10.3	0.01
Magnesium	mg/L	2.8	0.01
1,2-dichloropropane	mg/L	< 0.0001	0.0001
Chlorothalonil	mg/L	< 0.0001	0.0001
Propyzamide	mg/L	< 0.0001	0.0001
Chlornitrofen	mg/L	< 0.0001	0.0001
Simazine	mg/L	< 0.001	0.001
Thiobencarb	mg/L	< 0.0001	0.0001
Diazinon	mg/L	< 0.0001	0.0001
Isoxathion	mg/L	< 0.0001	0.0001
Fenitrothion	mg/L	< 0.0001	0.0001
EPN	mg/L	< 0.0001	0.0001
Dichlorvos	mg/L	< 0.0001	0.0001
Iprobenfos	mg/L	< 0.0001	0.0001
PCB	mg/L	< 0.0005	0.0005


Appendix 2

Analytical method and measured concentration of test item

1. Pretreatment of test solution

The test solution sampled was pretreated according to the flow scheme to prepare the sample for analysis.

Flow scheme



* Headspace method condition

Vessel: 125 mL vial container

Warming: 70°C, more than 20 min.

2. Method of analysis

The pretreated samples for analysis were quantitatively analyzed by gas chromatograph-mass spectrometry (GC-MS) under the following conditions to determine the concentration of the test item. The concentration of the test item in each sample for analysis was determined on the basis of a comparison of the peak area on the chromatogram of the sample with that of a standard sample. Some chromatograms obtained are shown in Appendix 3.

Analytical conditions

Instrument	Gas chromatograph-mass spectrometer
Gas chromatograph	Agilent 6890 Series Plus ⁺
Mass spectrometer	Agilent 5973N MSD

Gas chromatograph conditions

Column	HP-PONA film thickness 0.5 μ m (Agilent Technologies) 50 m \times 0.2 mm I.D. Fused silica
Column temperature	40°C(2 min) ^① \rightarrow 70°C(0 min) ^② \rightarrow 150°C(0.1 min)
Temp. rate	①15°C/min ②30°C/min
Carrier gas	Helium
Column flow	24.1 mL/min
Injection temp.	150°C
Injection volume	0.1 mL
Inlet mode	Split
Split ratio	13:1
Pressure	40 kPa

Mass spectrometer conditions


Ionization method	Electron ionization (EI)
Detecting method	Selected ion monitoring (SIM)
Measurement (m/z)	77
Ion source temp.	230°C
MS quadrupole temp.	150°C
Ionization voltage	69.9 eV
Interface temp.	200°C

3. Preparation of standard sample

The standard sample to determine the concentration of the test item in the sample for analysis was prepared as follows. The standard sample was prepared with correcting by the purity (99.8%) of the test item.

The test sample of 100.2 mg was precisely weighed with an electronic balance and dissolved in acetonitrile to obtain 1000 mg/L solution of the test item. The solution was diluted with acetonitrile to prepare 200 mg/L solution of the test item. And the solution was pretreated according to the flow scheme to prepare the 0.200 mg/L standard sample.

Flow scheme



4. Calibration curve

The test item solution of 20.0, 100, 200 and 400 mg/L were prepared by the same procedure as described in section 3. And they were pretreated according to the flow scheme of section 3 to prepare the standard sample of 0.0200, 0.100, 0.200 and 0.400 mg/L, respectively. These samples were analyzed according to the quantitative analytical conditions described in section 2. A calibration curve was drawn from the relationship between the concentrations of standard sample and the peak area on the chromatogram, and the determination was confirmed. The calibration curve is shown in Appendix 3. The determination limit of the test item was the lowest concentration of the standard sample (0.0200 mg/L) within the range of the calibration confirmed. Therefore, the determination limit of the test item in the test solution was 0.0200 mg/L in consideration of pretreatment.

5. Results of measurement

The results of the measured concentrations of the test item in the test solution are shown below.

Appendix table 2-1 Measured concentrations of test item in test solutions

Nominal concentration (mg/L)	Measured concentration (mg/L)								
	(Percentage of measured concentration versus that at each preparation %)								
	At the start	24 hours		48 hours		72 hours		At the end	Geometric mean
old		new	old	new	old	new			
Control	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
100	0.329	0.168 (51.1)	0.213	0.0880 (41.2)	0.0831	0.0666 (80.2)	0.0805	0.0768 (95.4)	0.117


old : before the renewal

new : after the renewal

n.d. : < 0.0200 mg/L

Appendix 3

Calibration curve and chromatogram



$$y = 8095886x$$

$$r = 0.998$$


Concentration (mg/L)	Peak area (-)
0.0200	159249
0.100	717033
0.200	1723007
0.400	3209712

Appendix figure 3-1 Calibration curve of 13F-OLE for analysis by GC-MS.

Standard sample 0.200 mg/L

データファイル : C:\MSDCHEM\1\DATA\94229\070820\0820S02.D N°イテ : 1
 測定日 : 20 Aug 2007 16:53 オペレータ :
 サンプル : Standard sample 0.200mg/L 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルア量 : 0.00
 MS積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231(B).M (ケミステーションインテグレータ)


94231



Control

データファイル : C:\MSDCHEM\1\DATA\94229\070820\31H0HZ.D N°イテ : 1
 測定日 : 20 Aug 2007 16:31 オペレータ :
 サンプル : 94231 本試験0h Control 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルア量 : 0.00
 MS積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231(B).M (ケミステーションインテグレータ)


94231



100 mg/L (Nominal concentration)

データファイル : C:\MSDCHEM\1\DATA\94229\070820\31H0HA.D N°イテ : 1
 測定日 : 20 Aug 2007 16:42 オペレータ :
 サンプル : 94231 本試験0h 100mg/L 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルア量 : 0.00
 MS積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231(B).M (ケミステーションインテグレータ)

94231




Appendix figure 3-2 GC-MS chromatograms at start of exposure.

Standard sample 0.200 mg/L

データファイル : C:\MSDCHEM\1\DATA\94229\070821\0821S01.D バイール : 1
 測定日 : 21 Aug 2007 14:56 オペレータ :
 サンプル : Standard sample 0.200mg/L 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルアmount : 0.00
 MS 積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231 (B).M (ケミステーション インテグレーション)


94231



Control

データファイル : C:\MSDCHEM\1\DATA\94229\070821\031H24HZO.D バイール : 1
 測定日 : 21 Aug 2007 15:07 オペレータ :
 サンプル : 94231 本試験 24h Control 換水前 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルアmount : 0.00
 MS 積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231 (B).M (ケミステーション インテグレーション)


94231



100 mg/L (Nominal concentration)

データファイル : C:\MSDCHEM\1\DATA\94229\070821\031H24HAO.D バイール : 1
 測定日 : 21 Aug 2007 15:18 オペレータ :
 サンプル : 94231 本試験 24h 100mg/L 換水前 装置 : Instrumen
 一般情報 : 希釈率 : 1.00
 サンプルアmount : 0.00
 MS 積分パラメータ : autoint1.e
 定量メソッド : C:\MSDCHEM\1\METHODS\94231 (B).M (ケミステーション インテグレーション)

94231



Appendix figure 3-3 GC-MS chromatograms at 24 hours after the exposure (before renewal).

Appendix 4

Solubility of test item in dilution water

1. Title

Solubility of test item in dilution water

2. Objective

The objective of this study is to estimate the solubility of the test item to dilution water.

3. Outline

Test item mixed with dilution water was stirred for 24 or 48 hours under the test temperature. After leaving at rest, the middle layer was sampled and analyzed.

4. Performance of test

4.1 Test equipments and instruments

Water bath :	Plastic tank Warming/cooling unit (Type HCA250, Sato craft)
Mixing apparatus :	Magnetic stirrer
Vessel :	Devised glass container (Interior volume : Approximately 600 mL)

4.2 Test conditions

(1) Test temperature :	$24 \pm 1^{\circ}\text{C}$
(2) Measurement :	Twice (24 and 48 hours after the mixture was stirred)
(3) Repitition :	24 hours n=3 (Sample-1, Sample-2 and Sample-3) 48 hours n=3 (Sample-4, Sample-5 and Sample-6)

4.3 Test procedures

- (1) Test sample and dilution water were mixed in a devised glass container to prepare approximately 100 mg/L* solution and sealed without headspace.

* The additive amount (38.5 μL) was calculated from the density of the test item (1.560 g/cm^3).

- (2) The test solution was stirred slowly using a magnetic stirrer under the test temperature in a water bath.
- (3) After the solution was stirred for 24 hours or 48 hours, the flask was settled in a water bath for approximately 1 hour.
- (4) After settling, middle layer was sampled and analyzed.

4.4 Analysis of test solution

- (1) Pretreatment of test solution

The middle layer of test solution was collected carefully from sampling spout of devised vessel by syringe. The collected solution was pretreated according to the flow scheme described in Appendix 2 1. Pretreatment of test solution.

- (2) Method for analysis

See Appendix 2 2. Method of analysis.

4.5 Preparation of standard sample

See Appendix 2 3. Preparation of standard sample

4.6 Calibration curve

See Appendix 2 4. Calibration curve.

5. Results

Measured solubility of the test item after 48 hours was higher than that of after 24 hours. Therefore, value of after 48 was adopted to the solubility in dilution water. The solubility of the test item to dilution water was 0.101 mg/L. The results of analyses are shown in below.

Appendix table 4-1 Value measured after stirring for 24 hours

Sample name	Measured value (mg/L)	Arithmetic mean (mg/L)
Sample-1	0.107	0.0927
Sample-2	0.0815	
Sample-3	0.0896	

Appendix table 4-2 Value measured after stirring for 48 hours

Sample name	Measured value (mg/L)	Arithmetic mean (mg/L)
Sample-4	0.104	0.101
Sample-5	0.0987	
Sample-6	0.101	

Additional data

Results of preliminary studies

1. Solubility of test item in dilution water

It was expected that the solubility of the test item in dilution water was below 100 mg/L, therefore, the measurement of the solubility of the test item in dilution water was conducted.

1) Preliminary study for measurement of solubility

(1) Method

Since the test item was expected to be volatile due to the chemical structure, the test item and the dilution water (dechlorinated tap water) were mixed and gently stirred in a devised glass container under closed system with no head space and test temperature ($24\pm 1^\circ\text{C}$) for 24 and 48 hours. For removal of insoluble substance, the procedure of gentle stirring and taking out from the middle layer of the settled solution for 1 hour was employed since centrifugation made the concentration of the test item decreased. The concentration of the test item in the collected sample was analyzed with pretreatment by gas chromatography - mass spectrometry (GC-MS). For 48 hours stirring, a sample of approximately 10 mg/L as nominal concentration was additionally measured.

(2) Result

Nominal concentration (mg/L)	Measured concentration (mg/L)	
	24-hour stirring hours	48-hour stirring
Approx. 100 (Sample-1)	0.0934	-
Approx. 100 (Sample-2)	0.129	-
Approx. 100 (Sample-3)	-	0.0949
Approx. 100 (Sample-4)	-	0.135
Approx. 10 (Sample-5)	-	0.0791

The measured value was around 0.1 mg/L with a little variance. In addition, the wide variance was not confirmed between approximately 10 and 100 mg/L as nominal concentration.

(3) Summary of preliminary study for measurement of solubility

From the results of the preliminary studies, the solubility of the test item in the dilution water was estimated at around 0.1 mg/L. Since centrifugation made the concentration of the test item decreased, the procedure of gentle stirring and taking out from the middle layer of the settled solution for approximately 1 hour was employed for removal of insoluble substance. Since the measured concentrations were almost same even in the different nominal concentrations of ten times (approximately 10 and 100 mg/L), it was thought that insoluble substance could be removed.

From the results mentioned above, in definitive study the devised glass container would be used for the preparation, and the procedure of gentle stirring and taking out from the middle layer of the settled solution for approximately 1 hour was employed in order to remove insoluble substance.

2. Effect on test organism

1) Preliminary study 1

(1) Method

After the test sample was added into the dilution water filled in Erlenmeyer flask with micro volumeter (Eppendorf Co., Ltd) to produce 100 mg/L as nominal concentration, the flask was immediately sealed with a plug not to produce head space. The solution was gently stirred by magnetic stirrer for approximately 48 hours to prepare the dispersed solution with suspended test item. This preliminary study was conducted by closed system without head space, since the test item was suspected to volatile. The test sample was employed in terms of volume using the density [1.560 g/cm³ (20°C)] for the preparation of test solution.

(2) Result

Nominal concentration (mg/L)	Left column : Cumulative mortality (%) Right column : Existence of abnormal response (abnormalities : *, no abnormalities : -)									
	3 hours		24 hours		48 hours		72 hours		96 hours	
Control	0	-	0	-	0	-	0	-	0	-
100	0	-	0	-	0	-	0	-	0	-

Type of test : Semi-static (renewal at every 24 hours)

Number of organisms/volume of test solution: Two fish/approx. 1 L

Aeration: Not conducted

No effect was observed on test organisms

2) Preliminary study 2

(1) Method

Prepare the dispersed solution with suspended test item by as same method preliminary study 1 was settled for approximately 1 hour under the $24\pm 1^{\circ}\text{C}$. The test solution was prepared by taking out from the middle layer of the settled solution. The stability study for the concentration of the test item was performed, while the test organisms were exposed to the test solution.

(2) Result

<Measured concentration of test item in test solution>

Nominal concentration (mg/L)	Measured concentration (mg/L) (percentage of measured concentration at start)	
	At the start	After 24 hours
Control	n.d.	n.d.
100	0.325	0.134 (41.2)

n.d. : <0.0200 mg/L

The saturation concentration of the test item in the test solution was decreased at 24 hours after the preparation.

(3) Summary of effect on test organisms (preliminary study)

The test item had no effect on the test organisms under the dispersed solution, which was prepared by mixing the test sample and the dilution water to produce 100 mg/L of an upper limit concentration on the test method by taking out from the middle layer of the solution. Since the test item was expected to volatile, the preliminary study was carried out under closed system. Since the test item concentration in the test solution decreased.

3. Operation of definitive study

1) Measurement of solubility of test item in dilution water

From the result of the preliminary study, the measurement of the solubility was performed using the solution taken out from the middle layer of the solution which was prepared by mixing the test sample and the dilution water to produce approximately 100 mg/L and stirred gently for 24 and 48 hours under the condition of $24\pm 1^{\circ}\text{C}$ and closed system. For removal of insoluble substance, the procedure of centrifugation or filtration was not used, but it of settling for approximately 1 hour after cease of stirring and then taking out from the middle layer of the settled solution was used as a method to remove as much as possible. The measurement of the test item concentration was employed for this test solution.

2) Definitive study

The definitive study was conducted as a limit test with a control and the saturated solution of the test item prepared by mixing the test item and dilution water to produce upper limit concentration (100 mg/L) of the test method, and stirring for approximately 24 hours. The definitive study was conducted by closed semi-static regime (renewal at every 24 hours). The test solution was prepared as follows; After the test sample was added, in terms of volume using the density, into the dilution water filled in Erlenmeyer flask with micro volumeter (Eppendorf Co., Ltd) to prepare test solution of 100 mg/L as nominal concentration, the flask was immediately sealed with a plug not to produce head space. The solution was gently stirred by magnetic stirrer for approximately 24 hours. After cease of stirring, the solution was settled for approximately 1 hour under $24\pm 1^{\circ}\text{C}$ and then test solution was prepared by taking out from the middle layer of the settled solution. The measurement of the test item in the test solution was carried out at the start of the exposure, before and after the renewal and at the end of the exposure.